







ORIGINAL ARTICLE

Awareness of chronic obstructive pulmonary disease and its risk factors among the adult population in the Jazan region, Saudi Arabia

Abdulrahman Hakami¹ , Mohammed H. Ghasham^{2*} , Khalid M. Alfaifi² ,
Abdulelah Almalki² , Shahad M. Alharbi² , Weam A. Mohammed², Raghd
A. Babaeer², Abdulrhman O. Alzhrani³ , Muhannad F. Alshrari⁴, Raghd M.
Alruwaili⁴

ABSTRACT

Background: Chronic obstructive pulmonary disease (COPD) is a leading cause of morbidity and mortality worldwide, yet public awareness remains limited, particularly in developing regions. This study aimed to assess COPD awareness and its associated risk factors among adults in the Jazan region, Saudi Arabia, and to identify factors associated with good knowledge.

Methods: A cross-sectional study was conducted among 513 adults using a structured questionnaire assessing sociodemographic characteristics, smoking behaviors, environmental exposures, and COPD-related knowledge. An awareness score was developed and categorized into low, moderate, and high levels. Associations were analyzed using chi-square tests and multivariate logistic regression.

Results: Only 31.4% had heard of COPD. While most identified smoking as a cause (468, 91.2%) and the lung as the affected organ (447, 87.1%), misconceptions were common: 266 (51.9%) believed COPD is rare, 187 (36.5%) thought it is curable with antibiotics, and only 192 (37.4%) recognized it as a lifelong disease. Most participants had moderate knowledge (358, 69.8%), followed by high (112, 21.8%) and low (43, 8.4%) levels. Cigarette smoking was significantly associated with lower knowledge (34.8% vs. 7.1%, $p < 0.001$). In multivariate analysis, male gender adjusted odds ratios (aOR = 0.612, $p = 0.037$) and lower income were associated with poorer knowledge, while higher income predicted better awareness (aOR = 1.329, $p = 0.004$).

Conclusion: COPD awareness in the Jazan region is suboptimal, with significant misconceptions and a disconnect between knowledge and behavior. Targeted public health interventions are needed to improve awareness and promote preventive practices.

Keywords: Chronic obstructive pulmonary disease (COPD), awareness, smoking, environmental exposure, socioeconomic factors, behavioral risk factors.

Introduction

Chronic obstructive pulmonary disease (COPD) is a progressive and preventable condition of the respiratory system, which is characterized by persistent limitation of airflow and chronic inflammation of the airways [1]. It is a major global health challenge as it is currently ranked among the top three leading causes of death worldwide [2]. According to the World Health Organization, it is responsible for approximately 3.2 million deaths annually, which accounts for nearly 5% of all deaths

Correspondence to: Mohammed H. Ghasham

*College of Medicine, Jazan University, Jazan, Saudi Arabia.

Email: Mohammed.ghasham0991@gmail.com

Full list of author information is available at the end of the article.

Received: 16 April 2026 | **Revised (1):** 03 May 2026 |

Accepted: 06 May 2026

45	worldwide [3]. The burden of COPD is expected to	Methods	102
46	increase further in the coming years due to aging of the	Study design and setting	103
47	populations, continued exposure to risk factors, and high	This study was a survey-based cross-sectional study	104
48	rate of underdiagnoses, particularly in low- and middle-	conducted over a 5-month period, from June 2024	105
49	income countries [4].	to November 2024. The study was carried out across	106
50	The most well-established risk factor for COPD is	multiple cities within the Jazan region, located in the	107
51	tobacco smoking, and is responsible for the majority	southern part of Saudi Arabia. The region comprises 13	108
52	of cases [5]. However, a significant proportion of	governorates with a population exceeding 1.5 million	109
53	patients develop COPD in the absence of smoking. This	people, distributed across urban and rural areas.	110
54	highlights the role of other important risk factors such as	Study population	111
55	exposure to biomass fuel, occupational dust, fumes, and	The study population included adult individuals who	112
56	environmental pollution [6]. The documented evidence	met the predefined inclusion and exclusion criteria.	113
57	has shown that up to 25%-45% of COPD cases occur	Eligible participants were mentally competent adults	114
58	among non-smokers, especially in regions where indoor	aged 18 years or older who were willing to participate	115
59	air pollution and occupational hazards are very common	in the study. Individuals under the age of 18, those	116
60	[7]. These findings emphasize that COPD is not solely a	who declined participation, and those with psychiatric	117
61	disease that occurs in smokers, but rather a multifactorial	illnesses or impaired judgment were excluded.	118
62	condition that is influenced by a range of environmental	Sampling and data collection procedure	119
63	and occupational exposures [8].	A random sampling technique was employed to recruit	120
64	Despite its high prevalence and preventable nature, the	participants, and the sample size was estimated to be	121
65	awareness of COPD among the general population is	400 individuals to ensure adequate statistical power.	122
66	very low. There are various studies conducted previously	Data were collected using a pretested and validated	123
67	in various regions, such as Saudi Arabia, which reported	questionnaire that had been adapted from previously	124
68	low awareness levels. The awareness of COPD among	published studies and translated into Arabic. Before the	125
69	the general population often ranged between 20% and	main data collection, a pilot study was conducted on 10-	126
70	40% [9,10]. Even among high-risk populations such as	15 participants to identify and resolve any ambiguities or	127
71	smokers, knowledge about COPD is limited. In addition,	technical issues in the questionnaire.	128
72	several misconceptions about COPD are present among	Questionnaire/study tool	129
73	the population among the population, with many	The questionnaire consisted of 23 items divided into	130
74	individuals believing that COPD is a rare disease, only	two main domains. The first domain included six	131
75	affects the elderly, or is curable with short-term treatment	questions addressing sociodemographic characteristics	132
76	[9]. Such misconceptions can delay diagnosis of this	and clinical information of the participants. The second	133
77	condition, reduce healthcare-seeking behavior, and	domain focused on assessing awareness of COPD	134
78	negatively impact the outcome of the disease.	and its associated risk factors, including knowledge	135
79	Importantly, awareness alone does not always lead to	of symptoms, risk factors, disease progression, and	136
80	appropriate changes in behavior. Some individuals may	complications. Each item in the awareness domain was	137
81	recognize smoking as harmful to their health yet continue	scored dichotomously, with a correct response assigned a	138
82	to engage in it [11]. This may also underestimate the risks	score of 1 and an incorrect or unknown response assigned	139
83	associated with passive smoking, occupational exposure,	a score of 0, yielding a total possible score of 17. Based	140
84	and indoor pollution. This disconnect between knowledge	on the total score, participants were categorized into three	141
85	and practice is an important barrier to effective disease	levels of awareness: low (0-5), moderate (6-11), and high	142
86	prevention [12]. Therefore, there is a need to improve	(12-17). This scoring system was newly developed for	143
87	the level of awareness, which must be accompanied	the purposes of this study, informed by the structure of	144
88	by strategies that promote behavioral change and risk	previously published tools assessing COPD awareness.	145
89	reduction.	Ethical approval	146
90	In Saudi Arabia, and particularly in the Jazan region, data	Ethical approval for this study was obtained from the	147
91	on COPD awareness and its determinants remain limited.	Standing Committee for Scientific Research at Jazan	148
92	Due to the unique sociodemographic characteristics and	University (reference number: REC-46/03/1184), dated	149
93	potential exposure to environmental and occupational	September 25, 2024. Participation was voluntary, and	150
94	risk factors in this region, it is very important to assess	informed consent was obtained from all participants	151
95	the level of public awareness and identify the existing	before data collection. Confidentiality and anonymity of	152
96	knowledge gaps in the region. Such information is crucial	the collected data were strictly maintained throughout the	153
97	for designing targeted public health interventions and	study.	154
98	educational programs tailored to the local population.		
99	Therefore, this study aims to assess the level of awareness		
100	of COPD and the risk factors associated with awareness		
101	among adults in the Jazan region, Saudi Arabia.		

155 **Statistical analysis**

156 Data analysis was performed using the Statistical Package
 157 for the Social Sciences (version 29.0; IBM Corp., Armonk,
 158 NY). Descriptive statistics were presented as frequencies
 159 and percentages for categorical variables, and as means ±
 160 standard deviations for continuous variables. An overall
 161 awareness score was calculated and categorized into
 162 low, moderate, and high levels. Associations between
 163 categorical variables and awareness levels were assessed
 164 using the chi-square test. Furthermore, multivariate
 165 logistic regression analysis was conducted to identify
 166 independent predictors of good awareness. Adjusted
 167 odds ratios (aORs) with 95% confidence intervals were
 168 reported, and a *p*-value of less than 0.05 was considered
 169 statistically significant.

170 **Results**

171 Our study included 513 participants, with a higher
 172 proportion of females (306, 59.6%) compared to males
 173 (207, 40.4%). Most participants were younger than 25
 174 years (211, 41.1%), with a mean age of 33.4 ± 12.7 years.
 175 Most participants had a bachelor's degree (332, 64.7%).
 176 Marital status was nearly balanced, with 259 (50.5%)
 177 single and 234 (93.6%) had children married. Among

178 married participants, 93.6% had children. Employment
 179 distribution was comparable across categories, with
 180 students (185, 36.1%), employees (184, 35.9%), and
 181 unemployed individuals (144, 28.1%). Slightly more
 182 participants resided in urban areas (281, 54.8%) than rural
 183 areas (232, 45.2%). The population was predominantly
 184 Saudi (499, 97.3%). Income levels were varied, with 159
 185 (31.0%) earning <5,000 SAR and 132 (25.7%) reporting
 186 >15,000 SAR (Table 1).

187 Figure 1 shows the distribution of employment sectors
 188 among participants. The majority were engaged in
 189 the educational sector (66.1%). Healthcare workers
 190 constituted 11.8%, followed by military personnel (8.6%).
 191 Smaller proportions were observed in industrial (3.6%)
 192 and freelance sectors (3.6%), while legal and financial
 193 services each accounted for 2.7%. The agricultural sector
 194 was minimally represented (0.9%).

195 Table 2 shows the distribution of smoking behaviors and
 196 environmental exposures among participants. Active
 197 smoking was relatively uncommon, with only 23 (4.5%)
 198 reporting shisha use and 23 (4.5%) cigarette smoking,
 199 while electronic cigarette use was reported by 11 (2.1%).
 200 In contrast, passive exposure was more frequent, with 116
 201 (22.6%) indicating exposure to smokers and 27 (5.3%)
 202 reporting time spent in shisha cafés. Environmental and


Table 1. Sociodemographic characteristics of study participants (N = 513).

203

Sociodemographic characteristics		Frequency N (%)
Gender	Female	306 (59.6)
	Male	207 (40.4)
Age (Years)	<25 years	211 (41.1)
	25-34 years	70 (13.6)
	35-44 years	101 (19.7)
	≥45 years	131 (25.5)
	Mean (SD)	33.4 (12.7)
	Range	18-70
Educational level	Secondary or less	152 (29.6)
	Bachelor's	332 (64.7)
	Postgraduate	29 (5.7)
Marital status	Single	259 (50.5)
	Married	234 (45.6)
	Widow/Divorced	20 (3.9)
Have children	Yes	234 (93.6)
	No	16 (6.4)
Job status	Unemployed	144 (28.1)
	Student	185 (36.1)
	Employee	184 (35.9)
Accommodation	City	281 (54.8)
	Village	232 (45.2)
Nationality	Saudi	499 (97.3)
	Non-Saudi	14 (2.7)
Family monthly income	<5,000 SAR	159 (31.0)
	5,000-9,999 SAR	103 (20.1)
	10,000-15,000 SAR	119 (23.2)
	>15,000 SAR	132 (25.7)

(N) Frequency, (%) Percentages.

footnote clarifying that these percentages apply to married participants only (n = 250), not the full sample (N = 513).



204 **Figure 1.** Distribution of different types of employment.

205 **Table 2.** Smoking behaviors, environmental and clinical exposure characteristics (N = 513).

		Frequency N (%)
Shisha smoking	No	490 (95.5)
	Yes	23 (4.5)
Cigarette smoking	No	490 (95.5)
	Yes	23 (4.5)
Electronic cigarette use	No	502 (97.9)
	Yes	11 (2.1)
Exposure to smokers (passive smoking)	No	397 (77.4)
	Yes	116 (22.6)
Time spent in shisha cafes	No	486 (94.7)
	Yes	27 (5.3)
Indoor biomass exposure (coal/wood)	No	489 (95.3)
	Yes	24 (4.7)
Occupational exposure (dust/fumes/chemicals)	No	447 (87.1)
	Yes	66 (12.9)
Diagnosed with COPD	No	508 (99.0)
	Yes	5 (1.0)

(N) Frequency, (%) Percentages.

206 occupational risk factors were less commonly reported,
 207 including indoor biomass exposure (24, 4.7%) and
 208 occupational exposure to dust, fumes, or chemicals (66,
 209 12.9%). Nearly all participants had not been diagnosed
 210 with COPD (508, 99.0%).

211 Table S1 shows the level of awareness and knowledge
 212 regarding COPD among participants. Only 161 (31.4%)
 213 had heard of COPD. However, most participants
 214 correctly identified the lung as the affected organ (447,
 215 87.1%) and recognized shortness of breath (373, 72.7%)

as a key symptom. A high proportion also acknowledged
 smoking as a cause (468, 91.2%) and the role of smoking
 cessation in prevention (484, 94.3%). Despite this,
 several critical misconceptions were evident. More than
 half perceived COPD as rare (266, 51.9%), and only
 192 (37.4%) correctly identified it as a lifelong disease.
 Additionally, 187 (36.5%) believed it could be cured with
 short-term antibiotics, and 299 (58.3%) did not recognize
 its potential to cause disability.

225 Figure 2 shows the distribution of COPD knowledge
226 levels among participants. The majority demonstrated
227 a moderate level of knowledge (358, 69.8%), while
228 112 (21.8%) had high knowledge, and only 43 (8.4%)
229 exhibited low knowledge.

230 Table 3 shows the association between sociodemographic
231 characteristics and COPD knowledge levels. Notably,
232 gender showed a significant association ($p = 0.010$),
233 with females having higher levels of good knowledge
234 (74, 24.2%) compared to males (38, 18.4%), while males
235 had a higher proportion of low knowledge (26, 12.6%).
236 Educational level was also significantly associated (p
237 $= 0.018$), with participants holding a bachelor's degree
238 demonstrating higher knowledge (86, 25.9%) compared
239 to other groups. Other variables, including age ($p =$
240 0.505), marital status ($p = 0.336$), having children (p
241 $= 0.320$), job status ($p = 0.072$), accommodation ($p =$
242 0.734), nationality ($p = 0.755$), and income ($p = 0.113$),
243 were not significantly associated with knowledge levels.

244 Table 4 shows the association between smoking
245 behaviors, environmental exposures, and COPD
246 knowledge levels. Cigarette smoking was the only factor
247 significantly associated with knowledge ($p < 0.001$),
248 with smokers demonstrating a substantially higher
249 proportion of low knowledge (8, 34.8%) compared to
250 non-smokers (35, 7.1%). In contrast, non-smokers had
251 higher proportions of moderate (348, 71.0%) and high
252 knowledge (107, 21.8%). Other variables did not show
253 significant associations.

254 Table 5 shows the multivariate logistic regression analysis
255 identifying predictors of good COPD knowledge. Gender
256 and family monthly income emerged as significant
257 predictors. Males had significantly lower odds of having
258 good knowledge compared to females (aOR = 0.612, $p =$
259 0.037). In contrast, higher family income was associated

with increased odds of good knowledge (aOR = 1.329, p
= 0.004). Age ($p = 0.190$), educational level ($p = 0.238$),
and marital status ($p = 0.820$) were not significantly
associated with knowledge after *adjustment*.

Discussion

COPD is a major global public health concern that
contributes significantly to morbidity and mortality
worldwide [13]. Despite its preventable nature, awareness
of COPD and its risk factors remains suboptimal,
especially in the developing countries of the world [14].
Therefore, understanding the level of public awareness
and associated behaviors is very important for designing
effective prevention strategies. This study aimed to assess
COPD awareness and its determinants among adults in
the Jazan region.

Notably, a key finding of this study is that there is
low overall awareness of COPD, with only 31.4% of
participants reporting that they had heard about COPD.
This finding aligned with findings from studies conducted
in Saudi Arabia, where awareness levels ranged between
20%-40%. For instance, a study by Al-Otaibi et al. [15]
reported limited recognition (20%) of COPD among the
general population. Another study by Oliveira et al. [16]
showed that only about a third of all the participants in the
study, 32.5%, know or have heard about COPD. Higher
awareness in Western studies may be due to stronger
health promotion campaigns and better integration of
respiratory education into primary care systems.

Despite limited overall awareness, participants
demonstrated good recognition of major risk factors,
particularly smoking, with 91.2% correctly identifying
it as a cause of COPD. This finding is consistent with
regional and international literature, where smoking
is widely recognized as a risk factor due to extensive




Figure 2. Knowledge level of participants about COPD.

295 **Table 3.** Association between sociodemographic characteristics and COPD knowledge levels (N = 513).

		COPD knowledge level			a Sig. Value
		Low N (%)	Moderate N (%)	High N (%)	
Age	<25 years	16 (7.6)	139 (65.9)	56 (26.5)	0.505
	25-34 years	7 (10.0)	51 (72.9)	12 (17.1)	
	35-44 years	9 (8.9)	71 (70.3)	21 (20.8)	
	≥45 years	11 (8.4)	97 (74.0)	23 (17.6)	
Gender	Female	17 (5.6)	215 (70.3)	74 (24.2)	0.010
	Male	26 (12.6)	143 (69.1)	38 (18.4)	
Educational level	Secondary or less	12 (7.9)	118 (77.6)	22 (14.5)	0.018
	Bachelor's	26 (7.8)	220 (66.3)	86 (25.9)	
	Postgraduate	5 (17.2)	20 (69.0)	4 (13.8)	
Marital status	Single	22 (8.5)	177 (68.3)	60 (23.2)	0.336
	Married	17 (7.3)	169 (72.2)	48 (20.5)	
	Widow/Divorced	4 (20.0)	12 (60.0)	4 (20.0)	
Have children	No	0 (0.0)	11 (68.8)	5 (31.3)	0.320
	Yes	20 (8.5)	167 (71.4)	47 (20.1)	
Job status	Unemployed	12 (8.3)	110 (76.4)	22 (15.3)	0.072
	Student	12 (6.5)	122 (65.9)	51 (27.6)	
	Employee	19 (10.3)	126 (68.5)	39 (21.2)	
Accommodation	City	23 (8.2)	193 (68.7)	65 (23.1)	0.734
	Village	20 (8.6)	165 (71.1)	47 (20.3)	
Nationality	Non-saudi	1 (7.1)	11 (78.6)	2 (14.3)	0.755
	Saudi	42 (8.4)	347 (69.5)	110 (22.0)	
Family monthly income	<5,000 SAR	13 (8.2)	122 (76.7)	24 (15.1)	0.113
	5,000-9,999 SAR	7 (6.8)	73 (70.9)	23 (22.3)	
	10,000-15,000 SAR	10 (8.4)	83 (69.7)	26 (21.8)	
	>15,000 SAR	13 (9.8)	80 (60.6)	39 (29.5)	

(a) Chi-Square Test.

296 **Table 4.** Association between smoking behaviors, environmental exposure, and COPD knowledge levels (N = 513).

		COPD knowledge level			a Sig. Value
		Low N (%)	Moderate N (%)	High N (%)	
Shisha smoking	No	38 (7.8)	344 (70.2)	108 (22.0)	0.060
	Yes	5 (21.7)	14 (60.9)	4 (17.4)	
Cigarette smoking	No	35 (7.1)	348 (71.0)	107 (21.8)	<0.001
	Yes	8 (34.8)	10 (43.5)	5 (21.7)	
Electronic cigarette use	No	41 (8.2)	349 (69.5)	112 (22.3)	0.137
	Yes	2 (18.2)	9 (81.8)	0 (0.0)	
Exposure to smokers (passive smoking)	No	29 (7.3)	276 (69.5)	92 (23.2)	0.143
	Yes	14 (12.1)	82 (70.7)	20 (17.2)	
Time spent in shisha cafes	No	38 (7.8)	340 (70.0)	108 (22.2)	0.124
	Yes	5 (18.5)	18 (66.7)	4 (14.8)	
Indoor biomass exposure (coal/wood)	No	42 (8.6)	339 (69.3)	108 (22.1)	0.560
	Yes	1 (4.2)	19 (79.2)	4 (16.7)	
Occupational exposure (dust/fumes/chemicals)	No	38 (8.5)	313 (70.0)	96 (21.5)	0.866
	Yes	5 (7.6)	45 (68.2)	16 (24.2)	
Diagnosed with COPD	No	43 (8.5)	353 (69.5)	112 (22.0)	0.335
	Yes	0 (0.0)	5 (100.0)	0 (0.0)	

(a) Chi-Square Test.

297 anti-smoking campaigns. Similarly, a study by Esam
 298 Mahmood et al. [14] reported that there is a moderate
 299 (37.4%) level of awareness of smoking as a key risk

factor for COPD and its worsening. De Fatima De 300
 Oliveira Graca et al. [17] show that 27.7% aware that 301
 quitting smoking is important for decreasing the severity 302

303 **Table 5.** Multivariate logistic regression analysis for predictors of good COPD knowledge.

	B	S.E.	Sig. Value	aOR	95% CI
Age	-0.018	0.014	0.190	0.982	0.957-1.009
Gender (Male)	-0.490	0.235	0.037	0.612	0.387-0.970
Higher educational level	0.253	0.215	0.238	1.288	0.846-1.962
Marital status (married)	0.066	0.289	0.820	1.068	0.606-1.882
Higher family monthly income	0.284	0.098	0.004	1.329	1.097-1.611
Constant	-1.780	0.522	0.001	0.169	—

304 of COPD symptoms. However, several critical gaps are
 305 present in our study, which include less recognized risk
 306 factors, such as passive smoking (22.6%), occupational
 307 exposure (12.9%), and indoor biomass exposure (4.7%).
 308 Similar gaps have been reported in studies from rural
 309 India and Iran, where environmental and occupational
 310 exposures are often underestimated. A study by Mishra
 311 et al. [18] shows that environmental and occupational
 312 air pollution is a major risk factor for COPD in the
 313 Indian subcontinent. Feizi et al. [19] show that ageing,
 314 occupational exposure to dust and fumes, indoor and
 315 outdoor air pollutants, genetic risk factors, and especially
 316 tobacco smoking are some of the risks. This discrepancy
 317 may be explained by the stronger emphasis of public
 318 health messaging on smoking, while indirect and
 319 environmental risks receive less attention.

320 Notably, another important observation is the presence
 321 of significant misconceptions regarding COPD. A
 322 considerable proportion of participants believed that
 323 COPD is rare (51.9%) or curable with short-term
 324 antibiotics (36.5%), and only 37.4% correctly identified
 325 it as a lifelong disease. These findings are consistent with
 326 studies conducted in Saudi Arabia, where misconceptions
 327 about chronic respiratory diseases are common.
 328 Similarly, Almuzaini et al. [9] show that 14.8% believed
 329 that COPD is curable with short-term antibiotics, which
 330 showed a key misconception. International studies have
 331 also reported similar misunderstandings, particularly in
 332 populations with limited exposure to health education
 333 [20]. These misconceptions may contribute to delayed
 334 diagnosis and poor disease management, ultimately
 335 increasing disease burden.

336 Furthermore, the distribution of knowledge levels
 337 further highlights that most participants had moderate
 338 knowledge (69.8%), with only 21.8% demonstrating
 339 high knowledge. This pattern has been observed
 340 in previous studies, which showed that while basic
 341 awareness exists, comprehensive understanding remains
 342 limited. Alsubaiei et al. [21] show that the mean \pm
 343 standard deviation knowledge score was 29.5 ± 4.2 out
 344 of 45 points, with most of participants falling into the
 345 moderate category (65.5%). The moderate knowledge
 346 of COPD often reflects fragmented information, where
 347 individuals recognize certain aspects (e.g., smoking risk)
 348 but lack a deeper understanding of disease progression
 349 and complications.

350 Notably, the association analysis revealed that female
 351 gender and higher education were linked to better
 352 knowledge in bivariate analysis. These are consistent

with findings from Saudi studies, which indicated that
 women and more educated individuals are generally more
 health-aware [22]. However, in the multivariate model,
 education lost its significance, while income remained a
 strong predictor. This revealed that socioeconomic status
 may play a more critical role than formal education
 alone. Similar findings have been reported in studies
 from Europe, where income influences access to health
 information, healthcare services, and overall health
 literacy [23].

Notably, one of the most striking findings of this study
 is the disconnect between awareness and behavior.
 Although awareness of smoking as a risk factor was
 high, some participants continued to smoke, and smokers
 demonstrated significantly lower knowledge levels.
 This inverse relationship has been reported in multiple
 studies, such as by Sikjær et al. [24], where smokers often
 underestimate personal risk despite general awareness.
 Additionally, continued exposure to passive smoking and
 occupational hazards further highlights that knowledge
 alone does not necessarily lead to behavior change.
 Cultural norms, social environments, and addiction
 likely contribute to this gap.

Public health implications

From a public health perspective, the findings of this study
 have several important implications. The coexistence of
 moderate awareness, persistent misconceptions, and risky
 behaviors showed that the current awareness strategies
 were insufficient. Without addressing these gaps, the
 burden of the COPD is likely to increase, especially in
 the regions with high exposure to environmental and
 occupational risks. Therefore, early prevention of the
 disease is very important as COPD shares risk factors
 with other chronic diseases which increase its overall
 health impact.

Limitations

There are several limitations of this study. The cross-
 sectional design of this study limits causal inference
 between awareness and behaviors. The data were
 self-reported, which may introduce recall and social
 desirability bias, particularly for the practice of smoking.
 The sample predominantly consisted of educated and
 urban individuals, which limited the generalizability
 of findings to rural or high-risk occupational groups.
 Additionally, the study did not assess behavioral
 outcomes such as physical activity or diet in depth.

399 *Future research directions*

400 Future research should include longitudinal designs, more
401 diverse populations, and objective measures of exposure.
402 Furthermore, interventional studies evaluating targeted
403 awareness programs and behavior change strategies are
404 needed in order to bridge the gap between knowledge
405 and practice.

406 **Conclusion**

407 This study shows that awareness of COPD in the
408 Jazan population remains insufficient. Several notable
409 misconceptions and gaps exist in the understanding of
410 the disease progression and risk factors. The knowledge
411 of smoking-related risks was high, but awareness of
412 environmental and indirect exposures was limited.
413 Factors such as gender and income influence knowledge
414 levels about COPD. Importantly, awareness did not
415 consistently translate into healthy behaviors. These
416 findings show the need for targeted, multifaceted
417 public health interventions to improve comprehensive
418 awareness and promote effective behavior change.

419 **List of Abbreviations**

420 aOR	Adjusted odds ratio
421 CI	Confidence interval
422 COPD	Chronic obstructive pulmonary disease
423 IBM	International Business Machines Corporation
424 KSA	Kingdom of Saudi Arabia
425 NY	New York
426 SAR	Saudi Arabian Riyal
427 SD	Standard deviation
428 SPSS	Statistical Package for the Social Sciences
429 WHO	World Health Organization

430 **Conflicts of interest**

431 The authors declare that they have no conflict of interest
432 regarding the publication of this article.

433 **Funding**

434 None.

435 **Consent to participate**

436 Informed consent was obtained from all participants
437 prior to data collection. Participation was voluntary, and
438 confidentiality and anonymity were strictly maintained, in
439 accordance with ethical approval granted by the Standing
440 Committee for Scientific Research at Jazan University
441 (reference number: REC-46/03/1184).

442 **Author details**

443 Abdulrahman Hakami¹, Mohammed H. Ghasham², Khalid M.
444 Alfaifi², Abdulelah Almalki², Shahad M. Alharbi², Weam A.
445 Mohammed², Raghd A. Babaeer², Abdulrhman O. Alzhrani³,
446 Muhannad F. Alshrari⁴, Raghd M. Alruwaili⁴

- 447 1. Department of Internal Medicine, College of Medicine,
448 Jazan University, Jazan, Saudi Arabia
- 449 2. College of Medicine, Jazan University, Jazan, Saudi Arabia
- 450 3. College of Medicine, Al Baha University, Al Bahah, Saudi
451 Arabia
- 452 4. College of Medicine, Al Jouf University, Sakaka, Saudi
453 Arabia
- 454

455 *Supplementary content (if any) is available online.*

References

- 456 1. Xu J, Zeng Q, Li S, Su Q, Fan H. Inflammation mechanism and
457 research progress of COPD. *Front Immunol.* 2024;2024:15.
458 <https://doi.org/10.3389/fimmu.2024.1404615>
459
- 460 2. Chapman KR, Mannino DM, Soriano JB, Vermeire PA,
461 Buist AS, Thun MJ, et al. Epidemiology and costs of
462 chronic obstructive pulmonary disease. *Eur Respir J.*
463 2006;27(1):188–207. [https://doi.org/10.1183/09031936](https://doi.org/10.1183/09031936.06.00024505)
464 .06.00024505
465
- 466 3. World Health Organization. Chronic obstructive pulmonary
467 disease (COPD) [Internet]. Geneva, Switzerland: WHO;
468 2023. Available from: [https://www.who.int/news-room/](https://www.who.int/news-room/fact-sheets/detail/chronic-obstructive-pulmonary-disease-(copd))
469 [fact-sheets/detail/chronic-obstructive-pulmonary-](https://www.who.int/news-room/fact-sheets/detail/chronic-obstructive-pulmonary-disease-(copd))
470 [disease-\(copd\)](https://www.who.int/news-room/fact-sheets/detail/chronic-obstructive-pulmonary-disease-(copd))
471
- 472 4. Boers E, Barrett M, Su JG, Benjafield AV, Sinha S, Kaye L, et
473 al. Global burden of chronic obstructive pulmonary disease
474 through 2050. *JAMA Netw Open.* 2023;6(12):e2346598.
475 <https://doi.org/10.1001/jamanetworkopen.2023.46598>
476
- 477 5. Lu W, Aarsand R, Schotte K, Han J, Lebedeva E, Tsoy E,
478 et al. Tobacco and COPD: presenting the World Health
479 Organization (WHO) tobacco knowledge summary. *Respir*
480 *Res.* 2024;25(1):338. [https://doi.org/10.1186/s12931-](https://doi.org/10.1186/s12931-024-02961-5)
481 [024-02961-5](https://doi.org/10.1186/s12931-024-02961-5)
482
- 483 6. Kc R, Shukla SD, Gautam SS, Hansbro PM, O’Toole RF. The
484 role of environmental exposure to non-cigarette smoke in
485 lung disease. *Clin Transl Med.* 2018;7(1):39. [https://doi.](https://doi.org/10.1186/s40169-018-0217-2)
486 [org/10.1186/s40169-018-0217-2](https://doi.org/10.1186/s40169-018-0217-2)
487
- 488 7. Salvi SS, Barnes PJ. Chronic obstructive pulmonary disease
489 in non-smokers. *Lancet.* 2009;374(9691):733–43. [https://](https://doi.org/10.1016/S0140-6736(09)61303-9)
490 [doi.org/10.1016/S0140-6736\(09\)61303-9](https://doi.org/10.1016/S0140-6736(09)61303-9)
491
- 492 8. Sama SR, Kriebel D, Gore RJ, Devries R, Rosiello R.
493 Environmental triggers of COPD symptoms: a case cross-
494 over study. *BMJ Open Respir Res.* 2017;4(1):179. [https://](https://doi.org/10.1136/bmjresp-2017-000179)
495 doi.org/10.1136/bmjresp-2017-000179
496
- 497 9. Almuzaini AS, Algeffari M, Alsohaibani A, Almutlaq LY,
498 Alwehaibi R, Almuzaini RA, et al. Awareness of chronic
499 obstructive pulmonary disease and its risk factors among
500 the adult population of the Qassim Region, Saudi Arabia.
501 *Cureus.* 2023;15(9):e44743. [https://doi.org/10.7759/](https://doi.org/10.7759/cureus.44743)
502 [cureus.44743](https://doi.org/10.7759/cureus.44743)
503
- 504 10. Al Bshabshe A, Al Shuqayfah N, Alahmari F, Alhomrany
505 A, Noreldin NM, Mousa WF, et al. Awareness of chronic
506 obstructive pulmonary disease (COPD) among the
507 general population in Aseer Region, Kingdom of Saudi
508 Arabia (KSA). *J Fam Med Prim Care.* 2023;12(6):1209–13.
509 https://doi.org/10.4103/jfmpc.jfmpc_2462_22
510
- 511 11. Wu J, Meng W, Ma Y, Zhao Z, Xiong R, Wang J, et al. Early
512 smoking lead to worse prognosis of COPD patients: a
513 real world study. *Respir Res.* 2024;25(1):140. [https://doi.](https://doi.org/10.1186/s12931-024-02760-y)
514 [org/10.1186/s12931-024-02760-y](https://doi.org/10.1186/s12931-024-02760-y)
515
- 516 12. Kahlke RM, McConnell MM, Wisener KM, Eva KW.
517 The disconnect between knowing and doing in health
518 professions education and practice. *Adv Health Sci*
519 *Educ Theory Pract.* 2020;25(1):227–40. [https://doi.](https://doi.org/10.1007/s10459-019-09886-5)
520 [org/10.1007/s10459-019-09886-5](https://doi.org/10.1007/s10459-019-09886-5)
521
- 522 13. Cao Z, Tong X, He L, Luo Y, Huang K, Li W, et al. Burden
523 of chronic obstructive pulmonary disease and its
524 attributable risk factors in 204 countries and territories,
525 1990–2021: results from the Global Burden of Disease
526 Study 2021. *BMJ Public Health.* 1990;4(1):2489. [https://](https://doi.org/10.1136/bmjph-2024-002489)
527 doi.org/10.1136/bmjph-2024-002489
528
- 529 14. Esam Mahmood S, Alqahtani AAT, Alghamdi BAA, Gazzan
530 MA, Alqahtani AMY, Alfaifi YNA, et al. Awareness of COPD
531

- 519 and its risk factors among the adult population of the
520 Aseer Region, Saudi Arabia. *Int J Chron Obstruct Pulmon*
521 *Dis.* 2023;18:23–35. [https://doi.org/10.2147/COPD.](https://doi.org/10.2147/COPD.S378064)
522 [S378064](https://doi.org/10.2147/COPD.S378064)
- 523 15. Al-Otaibi H, Homoud M, Alqarni M, Alsobhi G, Al-Otaibi G,
524 Alnahdi N. Level of awareness about chronic obstructive
525 pulmonary disease among active smokers. *Int J Chron*
526 *Obstruct Pulmon Dis.* 2026;21:580239. [https://doi.](https://doi.org/10.2147/COPD.S580239)
527 [org/10.2147/COPD.S580239](https://doi.org/10.2147/COPD.S580239)
- 528 16. Oliveira AS, Cortesão N, Gomes J, Pinto P, Shiang T, Alfaro T,
529 et al. Unmasking the hidden threat: cOPD awareness and
530 knowledge in Portugal. *Pulmonology.* 2024;30(3):207–9.
531 <https://doi.org/10.1016/j.pulmoe.2023.11.005>
- 532 17. Graca YDFDO, Yang L, Mingyu C, Alpona AB, Watanabe
533 T, Sawada Y, et al. Smoking attitudes, self-reported
534 practices, and COPD knowledge among adults aged 20-
535 59 years: insights from a Japanese sample. *Tob Induc Dis.*
536 2025;23:1–9. <https://doi.org/10.18332/tid/200855>
- 537 18. Mishra J, Acharya S, Taksande AB, Prasad R, Munjewar PK,
538 Wanjari MB. Occupational risks and chronic obstructive
539 pulmonary disease in the indian subcontinent: a
540 critical review. *Cureus.* 2023;15(6):e41149. [https://doi.](https://doi.org/10.7759/cureus.41149)
541 [org/10.7759/cureus.41149](https://doi.org/10.7759/cureus.41149)
- 542 19. Feizi H, Alizadeh M, Nejadghaderi SA, Noori M, Sullman
543 MJM, Ahmadian Heris J, et al. The burden of chronic
544 obstructive pulmonary disease and its attributable
545 risk factors in the Middle East and North Africa region,
546 1990-2019. *Respir Res.* 2022;23(1):319. [https://doi.](https://doi.org/10.1186/s12931-022-02242-z)
547 [org/10.1186/s12931-022-02242-z](https://doi.org/10.1186/s12931-022-02242-z) 548
- 549 20. Oweis J, Adams SG. Dispelling myths and misconceptions
550 about chronic obstructive pulmonary disease. *Med*
551 *Res Arch.* 2024;12(7). [https://doi.org/10.18103/mra.](https://doi.org/10.18103/mra.v12i7.5463)
552 [v12i7.5463](https://doi.org/10.18103/mra.v12i7.5463) 553
- 554 21. Alsubaiei ME, Frith PA, Cafarella PA, Quinn S, Al Moamary
555 MS, Mcevoy RD, et al. COPD care in Saudi Arabia:
556 physicians' awareness and knowledge of guidelines
557 and barriers to implementation. *Int J Tuberc Lung Dis.*
558 2017;21(5):592–5. <https://doi.org/10.5588/ijtld.16.0656> 559
- 560 22. Al-Khashan HI, Almulla NA, Galil SA, Rabbulnabi AA,
561 Mishriky AM. Gender differences in health education
562 needs and preferences of Saudis attending Riyadh
563 Military Hospital in the Kingdom of Saudi Arabia. *J*
564 *Fam Community Med.* 2012;19(3):172–7. [https://doi.](https://doi.org/10.4103/2230-8229.102317)
565 [org/10.4103/2230-8229.102317](https://doi.org/10.4103/2230-8229.102317) 566
- 567 23. Gero K, Backhaus-Hoven I, Höhmann A, Dragano N, Hoven
568 H. Low income and health literacy: a systematic scoping
569 review. *Arch Public Health.* 2025;83(1):291. [https://doi.](https://doi.org/10.1186/s13690-025-01781-3)
570 [org/10.1186/s13690-025-01781-3](https://doi.org/10.1186/s13690-025-01781-3) 571
- 572 24. Sikjær MG, Hilberg O, Fløe A, Dollerup J, Løkke A. Lack
573 of awareness towards smoking-related health risks,
574 symptoms related to COPD, and attitudinal factors
575 concerning smoking: an Internet-based survey conducted
576 in a random sample of the Danish general population. *Eur*
577 *Clin Respir J.* 2018;5(1):1506235. [https://doi.org/10.1080](https://doi.org/10.1080/20018525.2018.1506235)
578 [/20018525.2018.1506235](https://doi.org/10.1080/20018525.2018.1506235) 579